



Product Datasheet

Product Name	Interleukin-1 beta Porcine Recombinant
Cata No	CB500302
Source	<i>Escherichia Coli.</i>
Synonyms	Catabolin, Lymphocyte-activating factor (LAF), Endogenous Pyrogen (EP), Leukocyte Endogenous Mediator (LEM), Mononuclear Cell Factor (MCF), IL1F2, IL-1 beta.

Description

Interleukin-1b is produced by activated macrophages, IL-1B stimulates thymocyte proliferation by inducing il-2 release, b-cell maturation and proliferation, and fibroblast growth factor activity. IL1B proteins are involved in the inflammatory response, being identified as endogenous pyrogens, and are reported to stimulate the release of prostaglandin and collagenase from synovial cells.

Interleukin-1b Porcine Recombinant produced in E.Coli is a non-glycosylated, Polypeptide chain containing 154 amino acids and having a molecular mass of 17690 Dalton.

The IL-1b is purified by proprietary chromatographic techniques.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Biological Activity

The ED50 as determined by the dose-dependant stimulation of murine D10S cells is < 10 ng/ml.

Purity

Greater than 95.0% as determined by:

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.

Formulation

The protein was lyophilized with no additives.

Stability

Lyophilized Interleukin 1b although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL1b should be stored at 4°C between 2-7 days and for future use below -18°C.

For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles.

Sequence

The sequence of the first five N-terminal amino acids was determined and was found to be Met-Ala-Asn-Val-Gln.

*** For Non-Clinical Research Use Only ***